

Medizinische Fakultät
der
Universität Duisburg-Essen

Institut für Immunologie

Role of Viral Infection in Transplantation Medicine

In a u g u r a l - D i s s e r t a t i o n
zur
Erlangung des Doktorgrades der Medizin
durch die Medizinische Fakultät
der Universität Duisburg-Essen

Vorgelegt von
Asmae Gassa
aus Mülheim a. d. Ruhr
2016

Dekan: Herr Univ.-Prof. Dr. med. J. Buer
1. Gutachter: Herr Univ.-Prof. Dr. med. K. S. Lang
2. Gutachter: Herr Prof. Dr. med. O. Witzke
3. Gutachter: Frau Univ.-Prof. Dr. rer. nat. M. Sester, Saarland

Tag der mündlichen Prüfung: 22. Dezember 2016

Folgende Publikationen beinhalten Teile der Dissertation:

IL-10 induces T cell exhaustion during transplantation of virus infected hearts

Gassa A, Jian F, Kalkavan H, Duhan V, Honke N, Shaabani N, Friedrich SK, Dolff S, Wahlers T, Kribben A, Hardt C, Lang PA, Witzke O, Lang KS

Cellular Physiology and Biochemistry. 2016

High frequencies of anti-host reactive CD8+ T cells ignore non-hematopoietic antigen after bone marrow transplantation in a murine model

Gassa A, Kalkavan H, Jian F, Duhan V, Khairnar V, Shaabani N, Honke N, Carpinteiro A, Botezatu L, Crivello P, Dolff S, Ferencik S, Häussinger D, Khandanpour C, Fleischhauer K, Witzke O, Wahlers T, Hardt C, Lang PA, Lang KS

Cellular Physiology and Biochemistry. 2016

Contents

1 Introduction	7
1.1 The Adaptive Immune System	7
1.1.1 Antigen Recognition by T Cells.....	8
1.1.2 Major Histocompatibility Complex (MHC)	8
1.1.3 Cytokines.....	10
1.2 Lymphocytic Choriomeningitis Virus (LCMV)	11
1.2.1 Virus Biology	11
1.2.2 Immunopathology	12
1.2.3 T Cell Exhaustion	12
1.3 Tolerance in Transplantation Medicine	13
1.3.1 Bone Marrow Transplantation (BMT).....	13
1.3.2 Solid Organ Transplantation (SOT)	14
1.4 Challenges in Transplantation Medicine.....	15
1.4.1 Graft versus Host Disease (GvHD).....	15
1.4.2 Graft Rejection.....	16
1.4.3 Role of Viral Infection.....	17
1.5 Aims.....	18
2 Materials and Methods.....	19
2.1 Materials	19
2.1.1 Chemicals and Reagents	19
2.1.2 Media and Solution.....	19
2.1.3 Primer.....	20
2.1.4 Equipment	20
2.1.5 Software.....	21
2.1.6 Antibody	21
2.1.8 Mice	21
2.1.9 Lymphocytic Choriomeningitis Virus (LCMV).....	22
2.1.10 Statistical analysis	23
2.2 Methods	23
2.2.1 Bone Marrow Chimera.....	23
2.2.2 Heterotopic Heart Transplantation (HTX).....	24
2.2.3 Plaque Assay	24
2.2.4 Flow Cytometry	25

2.2.5 Depletion of CD8 ⁺ T Cells	26
2.2.6 Triggering of Innate Immunity	26
2.2.8 ELISA.....	26
2.2.9 RT-PCR	27
2.2.10 Analysis of Liver and Heart Enzymes	27
3 Results	27
3.1 Transgenic LCMV-GP is ubiquitously expressed	27
3.2 Host specific CD8 ⁺ T cells are not negatively selected after BMT	29
3.3 Introduction of LCMV-GP minor antigen in hematopoietic stem cells leads to deletion of host specific CD8 ⁺ T cells after BMT	32
3.4 Recipients ignore minor antigen (LCMV-GP) expressed in cardiac graft.....	33
3.5 Immunological tolerance in an LCMV model	34
3.5.1 LCMV carrier mice	35
3.5.2 LCMV transmitted by HTX leads to T cell exhaustion in WT mice.....	35
3.6 Breaking immunological tolerance	37
3.6.1 High frequencies of host specific CD8 ⁺ T cells induce limited GvHD after stimulation with LPS.....	38
3.6.2 IL-10 deficient mice die after transplantation of LCMV infected hearts.....	39
3.7 Graft rejection by memory T cells	41
3.7.1 Memory mice recipients reject LCMV infected heart transplant.....	42
4 Discussion	44
4.1 Host specific CD8 ⁺ T cells proliferate in presence of minor antigen in host after BMT.....	44
4.2 Triggering the innate immune system can lead to activation of host specific CD8 ⁺ T cells	45
4.3 Outlook for possible therapeutic methods in BMT	45
4.4 Immunized recipients have an increased risk for cardiac graft rejection	46
4.5 Role of LCMV infection in human beings	47
4.6 T cell exhaustion- a possible mechanism inducing tolerance?.....	48
5 Summary.....	50
6 References.....	51
7 Abbreviation.....	60

8 Tables and Figures63

9 Acknowledgment.....65

10 Curriculum vitae66

1 Introduction

The immune system is a powerful means against a broad spectrum of infectious agents in the environment, such as viruses, bacteria, fungi and prions. Successful elimination is guaranteed by two fractions which can interact intensely: On the one hand, the innate immune system provides granulocytes and macrophages that follow physiological barriers such as skin and mucous membrane. On the other hand, delayed but specific immune response is maintained by the adaptive immune system (Dempsey et al., 2003). Immune cells from both sections originate from hematopoietic stem cells in the bone marrow. A central capacity of the immune system is that it can distinguish between self and non-self-antigens (Jiang et al., 2009). In case of transplantation, a fully competent immune system recognizes foreign antigens on transplant tissue and in consequence, leads to rejection of the transplant. To avoid rejection of transplant tissue, suppression of the host immune system has to be maintained.

1.1 The Adaptive Immune System

The adaptive immune system differs from the innate immune system by its specific and delayed immune response. The adaptive or acquired immune system is also capable of memorizing experienced infections which lead to a fast and efficient elimination of invasive pathogens. Cellular components of this very specific immune response are antigen presenting cells (APCs), T cells and B- cells. Lymphocytes develop from hematopoietic stem cells and expose antigen specific receptors which recognize peptides and proteins. During maturation of lymphocytes somatic recombination of antigen specific receptors enables a highly specific B cell and T cell receptor repertoire. Theoretically, 10^{14} different B cell receptors (BCR) and 10^{18} different T cell receptors (TCR) are available (Lehrer, 2004). In general, two types of immune responses can be differentiated: the B cell derived humoral immune response and the T cell mediated cellular immune

response. Here in this study we focused on T cell mediated immune response and their interaction with foreign antigens.

1.1.1 Antigen Recognition by T Cells

T cells develop from common lymphoid progenitors (CLP) in the bone marrow and mature in the thymus. The TCR is a transmembrane protein, composed of two chains, the alpha and beta chains, which are associated with the CD3 complex (Borst et al., 1996; Call et al., 2005). Recognition of foreign antigens is only possible when peptides are exposed on the surface of the body's own cells. Foreign antigens can originate from viruses or intracellular bacteria which are processed intracellularly and then exposed on the surface; similarly for other pathogens or their fragments which are processed after endocytosis. T cells identify infected cells only if foreign peptides are presented on host specific glycoproteins which are defined as major histocompatibility complex (MHC) molecules. Discovery of highly polymorphic genes located in the MHC explained the powerful immune response to transplanted tissue (Murphy, K.P. (2012): Janeway's Immunobiology. 8th Ed. New York).

1.1.2 Major Histocompatibility Complex (MHC)

Characteristic for T cell recognition of antigens is the TCR which interacts with MHC surface molecules. MHC molecules are expressed on lymphoid and non-lymphoid tissue except on erythrocytes. There are two types of MHC: MHC class I and MHC class II. Their function is to present foreign antigens on the cell surface to T cells so that viral and / or bacterial replication in cells can be stopped by cell killing. Peptides derived from the cytosol are transported to the endoplasmic reticulum where they fuse with newly synthesized MHC class I molecules. This complex is exposed on the cell surface of APCs which present foreign antigens to CD8⁺ T cells. Peptides generated by degradation of proteins in intracellular endosomal vesicles combine with MHC class II molecules. CD4⁺ T cells then

recognize the complex of peptide and MHC class II molecules on the cell surface of APCs. CD8⁺ T cells recognize antigen in the context of MHC class I molecules whereas CD4⁺ T cells recognize antigen in the context of MHC class II, a process which is called MHC-restriction. CD4⁺ and CD8⁺ T cells recognize foreign antigen on self-MHC molecules (self MHC-restricted) or self or foreign antigen on foreign-MHC molecules (allo MHC-restricted). Genetically, MHC molecules are encoded on chromosome 6 in humans and on chromosome 17 in mice and are highly polymorphic which makes it difficult for viruses and bacteria to evade the immune response ("Complete sequence and gene map of a human major histocompatibility complex. The MHC sequencing consortium," 1999).




Figure 1.1 Genetic organization of the major histocompatibility complex in human and in mouse.

(Murphy, K.P. (2012): Janeway's Immunobiology. 8th Ed. New York. Chapter 6)

1.1.3 Cytokines

Cytokines are key molecules which enable cells to communicate and to interact via cytokine secretion. Cytokines are small proteins that have specific impact on target cells. Cytokines vary in their names depending on their origins. Lymphokines, are cytokines made by lymphocytes, monokines come from monocytes, chemokines are named due to their chemotactic activity and interleukins are cytokines which are made by leucocytes and serve as cell interacting molecules between leucocytes (Moreno Brea et al., 2009). Regulation of suppressive or activating effects in target cells is maintained by pro-inflammatory and anti-inflammatory cytokines. In this study, interferon- γ played a crucial role as pro-inflammatory and interleukin-10 as inhibitory immunosuppressive cytokine.

1.1.3.1 Interferon- γ (IFN- γ)

IFN- γ is one of the different types of interferon besides IFN- α and IFN- β . IFN- γ is also known as type-II interferon, whereas IFN- α and IFN- β belong to type-I interferon. It activates innate immune responses and mediates anti-viral effects. During viral infection, the main sources of IFN- γ production are CD8⁺ and CD4⁺ T cells. Signal transduction undergoes the Jak-Stat pathway. IFN- γ mediates activation of macrophages and granulocytes for phagocytosis and cell proliferation (Schroder et al., 2004).

1.1.3.2 Interleukin-10 (IL-10)

IL-10 is an immunoregulatory cytokine that is associated with T cell exhaustion. Its function during viral infection is described to be immunosuppressive and anti-inflammatory by suppressing cytokine production and proliferation of CD8⁺ and CD4⁺ T cells, leading to viral persistence. It inhibits NF- κ B activation through ill-defined mechanisms. A variety of cells can produce IL-10 such as T cells, B cells and APCs (Blackburn et al., 2007; Moore et al., 2001).

1.2 Lymphocytic Choriomeningitis Virus (LCMV)

LCMV has become an indispensable model in immunology. Due to viral infection experiments in mice, it was possible to start to understand T cell response and MHC-restriction (Zinkernagel et al., 1979, 1997).

1.2.1 Virus Biology

LCMV is an arenavirus, which is a non-cytopathic virus. Regarding the genome, LCMV contains a single stranded RNA. Its physiological reservoir is the house mouse and hamster (Childs et al., 1992). The first arenavirus, LCMV, has been isolated by Charles Armstrong in 1933. During an epidemic study in St. Louis, the virus has been detected. As common house mice provide an LCMV reservoir, human beings can be in contact with the virus. In our days, contamination with LCMV in the population is approximately 5%. Infection with LCMV in humans can be asymptomatic but can also provoke meningitis. In immunosuppressed patients such as transplanted recipients, donor derived LCMV infection in solid organ transplantation (SOT) leads to higher morbidity and mortality. Vertical transmission of LCMV in pregnancy is not well studied but first approach assumes complications and higher morbidity for the fetus.

To get insight into mechanism in viral clearance, different strains of virus exist (Ahmed et al., 1988; Zinkernagel et al., 1986). Concerning the immune response, we distinguish between strains inducing a chronic and an acute infection. Strains as Armstrong and WE enable an acute infection in mice, the virus can be cleared 8 days *post-infection*. Armstrong is a neurotropic LCMV strain whereas WE is a visceral strain. Chronic infections are well described for Clone 13 and docile. Looking closer at the infection with strain WE of LCMV, CD8⁺ T cell response can be well characterized. Great achievements in immunological understanding of T cell response during viral infection has been done with LCMV. In this study, acute infection in transplantation medicine was the focus. Therefore, LCMV-WE has been used for acute infection in mice.

1.2.2 Immunopathology

LCMV infection is a very good model to study and understand immunopathology. During viral infection, antigen specific CD8⁺ T cells kill infected cells. During acute infection with LCMV strain WE high dose (2×10^6 PFU), T cell mediated tissue damage in liver takes place. Virus can be cleared but at the same time viral infected cells get eliminated. In consequence, tissue damage marker for liver are elevated. Liver enzymes such as alanine transferase (ALT) and bilirubin are elevated. Lactate dehydrogenase (LDH), an unspecific cell damage marker, is massively increased. Chronic infection with LCMV can lead to constant activation of T cells which consequently lead to massive immunopathology and death. In this study, virus titer was determined in organs to look for viral persistence and ALT and LDH were measured in sera (K. S. Lang, Recher, Navarini, et al., 2005; P. A. Lang et al., 2010).

1.2.3 T Cell Exhaustion

T cell exhaustion is defined as a loss of effector function (Wherry, 2011; Yi et al., 2010). Gradual loss can be distinguished in stages which are determined by diminished or vanished production of effector cytokines such as IFN- γ , increased expression of inhibitory receptors and deletion of antigen-specific T cells. Furthermore T cell exhaustion correlates with viral persistence (Wherry et al., 2007).




Figure 1.2 T cell exhaustion and immunopathology (P. A. Lang et al., 2010)

1.3 Tolerance in Transplantation Medicine

Immune tolerance is defined as a state of specific immunological unresponsiveness in the immunocompetent host (Dong et al., 1999). Transplantation medicine has improved in the last decades and since the use of new immunosuppressive drugs. Still, long-term survival and chronic rejection remain major challenges in transplantation medicine. Moreover, immunosuppressive drugs bring along many side effects such as toxicity, malignancies and cardiovascular disease (Mihatsch et al., 1998). Therefore, understanding tolerance mechanism in transplantation medicine is crucial.

1.3.1 Bone Marrow Transplantation (BMT)

Hematopoietic stem cell transplantation (HST) is an important therapy for patients with acute myeloid leukemia, myelodysplastic syndrome, refractory relapsed lymphoma and very severe aplastic anemia. HST is the transplantation of multipotent hematopoietic stem cells from bone marrow, peripheral blood, or umbilical cord blood. It can be autologous (patients' own stem cells) or allogenic (donor stem cells). Allogenic stem cell transplantation requires that the transplant carries compatible transplantation antigens which are determined by the variability of three or more loci of the MHC complex. In humans, these MHC class I and class II loci are named human leucocyte antigens (HLA) class I and class II. Once HLA testing is assessed, compatible donors are selected for HST. Compatibility is verified by the two major categories of HLA type (type I and type II). The risk that the transplanted "new" immune system attacks the recipient tissue increases with every HLA-mismatch and causes severe illness. This phenomenon is called graft versus host disease (GvHD). The risk of GvHD increases with mismatches of HLA Type-I genes. Still, fully HLA-matched donor and recipients as in the case of syngeneic HST (siblings) can lead to GvHD.

1.3.2 Solid Organ Transplantation (SOT)

Due to the discovery of HLA in 1967 (Turner, 2004), transplantation medicine progressed enormously. Solid organ transplantation means transplantation of organs and tissues from deceased or living donors. Organs that can be transplanted are the heart, kidney, liver, pancreas, intestine and thymus. Considering tissue transplantation, skin, heart valves, cornea, bones and tendons can be transplanted. Autografts are autologous transplantation within the same person e.g. skin-transplantation. In case of allograft transplantation, certain conditions have to be fulfilled. First of all, as HLA mismatch correlates with allograft rejection, HLA matching between donor and recipient has to be determined. This is assessed by serologic or DNA typing methods.

Furthermore, detection of alloreactive T cells opposed to recipient is measured in mixed lymphocyte reaction (MLR). Histoincompatibility can be verified *in vitro*. If HLA match is determined by DNA typing but MLR reveals histoincompatibility, the donor is not compatible for the recipient. As results usually take a long time to be ready and organ availability is short, HLA match is subordinated to organ demand. Immunosuppression then supports allograft survival by suppressing the immune response nonspecifically. Similar to alloreactive T cells, alloreactive antibodies can be identified in donor-recipient serologic cross matching. This analysis is particularly important for vascularized organs such as kidney and heart. Presence of such antibodies induce hyperacute rejection (Kissmeyer-Nielsen et al., 1966).

1.3.2.1 Heart Transplantation (HTX)

Heart transplantation is the therapy of choice for patients with heart insufficiency. So far, limited amount of available donor organs and the rejection of hearts after transplantation are major problems in heart transplantation. The first heart transplantation was performed in South Africa in 1967. Over the last two decades, selection methods for donor and recipients have increasingly improved. In addition, introduction of immunosuppressive drugs such as cyclosporine

opened up new opportunities in transplantation medicine. Overall, after transplantation, survival in the first year is more than 90% and in the second year approximately 70%. The field of heart transplantation is constantly progressing. Advances in organ preservation, immune monitoring and immunosuppressive regimens are likely to lead to a better outcome and to advances in the quality and the length of life of heart transplant recipients (Hoffman, 2005; Hunt et al., 2008).

1.4 Challenges in Transplantation Medicine

In the last 20 years, transplantation medicine has strongly improved thanks to immunosuppressive drugs, better surgical techniques and controlling infectious disease. Still, lethal problems exist in transplantation medicine and tolerance mechanisms are not understood yet.

1.4.1 Graft versus Host Disease (GvHD)

Still limitation to successful bone marrow transplantation (BMT) comes from severe GvHD, which occurs in up to 40% - 60% of cases (Jacobsohn et al., 2007; Jagasia et al., 2012) and leads to 15% of deaths after HST. GvHD is clinically grouped in an acute and chronic GvHD defined by the time of GvHD onset after HST with a cutoff of 100 days. Mechanistically chronic GvHD (cGvHD) usually involves several factors including B cell activation, production of autoantibodies and absence of T-regulatory cells (Treg) (Schroeder et al., 2011).

Acute GvHD (aGvHD) is characterized by the reaction of CD8⁺ T cells which are activated by antigens presented on host MHC class I molecules (Ferrara et al., 1991). Therefore, T cell depletion of the transplant prior BM transfer is performed in cases of HLA mismatch or unrelated donors and significantly reduces aGvHD (Ho et al., 2001; Kroger et al., 2002; Remberger et al., 2001). Due to the high number of registered bone marrow donors most of BMT are done in complete MHC class I match. Nevertheless, around 40% of patients develop aGvHD despite complete HLA-matching (Ferrara et al., 2009) and T cell depletion (Champlin et al.,

2000). Preparative regimens are critical in terms of treatment of the primary disease and as prevention of graft rejection. Common conditioning regimens are either myeloablative or non-myeloablative defined by the pancytopenia caused. During the last 20 years, an alternative preparative regimen has been developed. This therapy is called reduced intensity conditioning (RIC) and causes less tissue damage and is associated with less GvHD (Mielcarek et al., 2003). RIC as an intermediate category of regimens is commonly used for elderly people to improve the non-relapse mortality by reducing toxicity of treatment by either whole body irradiation or chemotherapy (Abdul Wahid et al., 2014; Mohty et al., 2010).

1.4.2 Graft Rejection

Allograft rejection is a major complication in solid organ transplantation. It is defined as a destructive immune response against foreign tissue or organ which can be mediated via cellular or humoral immune response. Three categories of allograft rejection can be distinguished according to their mechanism and time of occurrence conferring to guidelines of the international society of heart and lung transplantation (Task Force 2: Immunosuppression and Rejection). Hyperacute allograft rejection happens within minutes or hours after transplantation. It is mediated via preformed antibodies hostile to HLA type I molecules expressed on endothelium. It leads to immediate loss of an allograft.

Further, acute cellular rejection occurs within 6 months in case of HTX. As the name itself explains, mainly T cells are responsible for infiltration into the allograft. APCs of immunocompetent host migrate from foreign donor organ to lymphoid tissue and prime T cells. This in consequence leads to direct allorecognition. Recipients' APCs are also capable to present peptide of allograft to T cells by indirect allorecognition (Davis et al., 2004).

Chronic allograft rejection limits long-term survival of transplant recipients. Beyond the 2nd year after HTX, cardiac graft recipients suffer from cardiac allograft vasculopathy (CAV) in 30% of cases. Chronic rejection is multifactorial. Viral infection or reactivation such as cytomegalovirus (CMV) can cause chronic rejection. Graft rejection can be well determined by biopsy. Heart specific tissue

damage markers such as creatinine kinase-MB (CK-MB) and troponine I (Top I) can give a hint (Dengler et al., 1998). Treatment is very often too late and has to be optimized.

1.4.3 Role of Viral Infection

The main reason for graft rejection is viral infection during heart transplantation. 40% of the heart rejections in humans are associated with infections (Jung et al., 2011; Yusen et al., 2014). Those infections are mainly caused by herpes viruses. Considering viral infection in transplantation, we distinguish between reactivation of latent infections in donor organs or recipients and unexpected transmission of viral infections through organ transplantation which is defined as donor-derived viral infection. Even though, transmission of acute or latent infection to organ transplant recipients occurs only in about 0.2% of transplantation. Once it occurs, it leads to significantly increased morbidity and mortality in the recipients (Grossi et al., 2009). In transplantation medicine, expected viral infections detected by serologic analysis are known for CMV, hepatitis B virus (HBV) and Epstein-Barr-virus (Razonable, 2011). Special prophylaxis and infection management is preserved in this setting. Of greater concern are the unexpected viral infections transmitted via donor organs. Several reports of infectious diseases through SOT are published and estimated number of unreported cases has to be considered. Donor-derived viral infections include a growing number of pathogens such as HBV, herpes viruses, human T- cell lymphotropic viruses (HTLV) 1 and 2, West Nile virus, Rabies, LCMV, polyomavirus, parvovirus B19 and many other viruses. In most of such transmissions the mortality is high (Ison et al., 2011; Srinivasan et al., 2005; Winston et al., 2014). Detection of infectious disease in donor organs is limited by technical means and by the short time period between death of donor and the use of transplant. So, routinely evaluation of donors for infectious disease is determined by antibody detection. Nevertheless, a time window of false negative detection for infectious disease exists. For example, for hepatitis C virus (HCV) the window from infection to seroconversion is 30-70 days. Improvements have been

done by nucleic acid testing (NAT), so that the window could be reduced. But still, some infections remain undetected (Fishman, 2007; Fishman et al., 2009).

1.5 Aims

Transplantation medicine is one of the most challenging fields in medicine. Mechanisms establishing tolerance towards graft and host respectively in an immunocompetent environment has not been understood yet. Factors influencing tolerance can be on the one hand minor antigen mismatches despite full HLA-matched donor and recipient. Another problem is viral infection during BMT and SOT which emerged being life-threatening complications. In this study reported here, LCMV was a very good model to analyze differences of immune responses towards minor antigen and virus replication during murine bone marrow and heterotopic heart transplantation. The aim of this study was to investigate the interaction between LCMV specific CD8⁺ T cells and presentation of LCMV peptide or virus. We hypothesized that only presence of replicating virus can break immunological tolerance.

2 Materials and Methods

2.1 Materials

2.1.1 Chemicals and Reagents

Chemicals were obtained from AppliChem (Darmstadt), Merck (Darmstadt), Roth (Karlsruhe) or Sigma-Aldrich (Munich).

Carboxyfluorescein Succinimidyl Ester (CFSE)	Invitrogen, Darmstadt
Fluorescence Activating Cell Sorter (FACS)	BD Bioscience, Heidelberg
Lysing Solution	BD Bioscience, Heidelberg
Fetal Calf Serum (FCS)	Biochrom AG, Berlin
Isofluran	Delta Select GmbH, Pfullingen
Lipopolysaccharides of <i>Escherichia coli</i> (LPS)	Sigma, Munich
TriFast®TRIZol	PeqLab, Erlangen
Mouse IL-10 ELISA Ready-SET-Go!®	eBioscience
QuantiTect Reverse Transcription Kit	QIAGEN, Hilden
SYBR®Green	(AppliedBiosystems, Darmstadt)

2.1.2 Media and Solution

Dulbecco's Modified Eagle's Medium (DMEM)	PAN Biotech GmbH, Aldenbach
DPBS w/o Mg ²⁺ , Ca ²⁺	PAN Biotech GmbH, Aldenbach
Phosphate Buffered Saline (PBS)	DPBS, ddH ₂ O
FACS-Buffer	PBS 2% (v/v) FCS 0.1% (v/v) NaN ₃ 2 mM EDTA

Erythrocyte-Lysis-Buffer	0.5 M NH ₄ Cl 10mM KHCO ₃ 0.1mM EDTA pH 7.42
EDTA-Solution	0.5 M EDTA pH 8.0
Iscove's Modified Dulbecco's Medium (IMDM)	175 ml dH ₂ O, 50 ml 9xIMDM, 2-4 ml 2M NaHO, sterile filtration, 5% (v/v) FCS, 2% (w/v) PSG
<i>Overlay</i> (Plaque Assay)	50% (v/v) IMDM Solution 50% (v/v) 2% Methylcellulose

2.1.3 Primer

qPCR-WEGPforw (LCMV-GP)	CGA GCA TCA AAG CTG TGT ACA AT
qPCR-WEGPrev (LCMV-GP)	AAA AGG AAG CTG ACC AGT GCT AA
GAPDH	Mm_Gapdh_3_SG QuantiTect Primer Assay (200), Qiagen GmbH, Hilden

2.1.4 Equipment

ELISA Reader	AnthosLabtecInstruments GmbH, Wals-Siezenheim
FACSCalibur	BD Bioscience, Heidelberg
Fluoreszenz-Mikroskop HS BZ-9000	Keyence GmbH, Neu-Isenburg
Kryostat CM 3050S	Leica, Wetzlar
NanoDrop ND-1000	Peqlab Biotechnologie GmbH, Erlangen
TissueLyserII	QIAGEN, Hilden
Micro 220R-Centrifuge	Andreas Hettich GmbH, Tuttlingen

2.1.5 Software

FacsDiva v6.2.1	BD Bioscience
FlowJo 7.6.1	TreeStar Inc., Ashland (OR, USA)
GraphPad Prism5	GraphPad Software, La Jolla (CA, USA)
MicrosoftOffice 2007	MicrosoftCorporation, Redmont (WA, USA)

2.1.6 Antibody

In this study, antibodies were directly coupled with fluorescence. Antibodies were obtained from BD Pharmigen (Heidelberg) and eBioscience (Frankfurt). Fluorochromes used were FITC, PE, PerCP, APC, PeCy7, AmCyan and Pacific Blue.

2.1.8 Mice

All mice used in these studies were maintained on the C57BL/6 (B6) genetic background. All animals were housed in single ventilated cages under standard conditions. Animal experiments were authorized by the Landesamt für Natur, Umwelt und Verbraucherschutz in Nordrhein Westfalen (Recklinghausen, Germany) and performed in accordance with the German law for animal protection. During survival experiments mice which showed a severe phenotype with weight loss (>20% of starting weight), reduced movement, and general sickness have been killed at indicated time points assuming time of death. Besides the C57BL/6 (B6) mice which were used as wild type (WT) mice different mouse strains have been used in these experiments.

DEE mice

DEE mice are transgenic mice that express ubiquitously the glycoprotein of LCMV under the H2k promoter (Hunziker et al., 2003). On the one hand, these mice

served as recipients for the generation of bone marrow chimera mice and on the other hand, the *DEE* hearts were used as donors for the heterotopic heart transplantation.

IL-10^{-/-} mice

IL-10 deficient mice lack secretion of the immune modulatory cytokine IL-10 (Kuhn et al., 1993). As recipient they received a heterotopic transplanted heart.

P14 mice

P14 T cells recognize the LCMV-GP on MHC class I. The P14 mice express an LCMV-GP33-41 specific TCR on CD45.1 positive CD8⁺ T cells (Pircher et al., 1989) and were used for adoptive transfer experiments and as bone marrow donors.

Sv/129 mice

Sv/129 mice have a different background from C57BL/6 mice. They differ in gene loci and are used as two mice strains with different backgrounds for e.g. stroke research (Fujii et al., 1997; Gerlai, 1996).

Tcrb^{-/-} mice

Tcrb^{-/-} mice are genetically engineered immunodeficient mice which lack the T cell receptor beta chain (Funabashi et al., 2001). After intravenous injection of LCMV-WE, the virus persisted in the mice. Their hearts were used as LCMV carrier donor hearts for heterotopic heart transplantation.

2.1.9 Lymphocytic Choriomeningitis Virus (LCMV)

LCMV strain WE originally obtained from F. Lehmann-Grube (Heinrich Pette Institute, Hamburg, Germany) was propagated in L929 cells. Mice were infected intravenously in the tail vein. Applied concentration was either 200 plaque-forming units (PFU) or 2x10⁶ PFU. Viral titers were measured in Plaque Assay using mouse fibrosarcoma cell line derived from C57BL/6 mice (MC57 cells).

2.1.10 Statistical analysis

If not differently stated data are expressed as means and S.E.M. Student's t-test was used to detect statistically significant differences between groups. Significant differences between several groups were detected by two-way analysis of variance (ANOVA). The level of statistical significance was set at * $P < 0.05$, ** $P < 0.01$ and *** $P < 0.001$.

2.2 Methods

2.2.1 Bone Marrow Chimera

Creating bone marrow chimera mice is a very useful method to get further insights into immunological processes such as influence of donor derived hematopoietic stem cells on immune responses. Especially, in the field of transplantation medicine, bone marrow chimera experiments play a crucial role for tolerance model (Umemura et al., 2001). This model makes it possible to distinguish between effects on the immune system in recipient induced by cells coming from the donor derived hematopoietic or non-hematopoietic system.

2.2.1.1 Preparation of Donor BM

For the preparation of donor bone marrow, one mouse had to be killed to gain bone marrow for transplantation of six mice. Donor and recipients were sex-matched. After disinfecting the dead mouse, the abdominal skin was cut open to reach the legs till the hip. Femur and tibia got totally removed without breaking the bones. Forceps and scissors were necessary to fulfill the procedure. When the bones were ready to be flushed, they had to be cleaned in 70% ethanol for three minutes and washed with PBS. With help of a 21-gauge needle and medium, the bone marrow got flushed after cutting open both sides of each bone. The bone marrow (which looked like red, long, thin tissue) had to be collected in a 50 ml

tube and vortexed into a single-cell suspension to passage through a strainer. Re-suspended in medium, the bone marrow was ready for injection. Around 50×10^6 bone marrow cells had to be injected intravenously in the tail vein of each mouse (Holl, 2013).

2.2.1.2 Recipient Preparation

Six to eight weeks old recipients underwent a full body irradiation by X-ray with 9.5 Gy. 24 hours after irradiation, mice received sex-matched bone marrow diluted in medium by tail-vein injection. 30 days after bone marrow transplantation reconstitution of bone marrow took place in the host.

2.2.2 Heterotopic Heart Transplantation (HTX)

Heart allografts were vascularly anastomosed in an intraabdominal location using the technique described previously (Corry et al., 1973; Wu et al., 2006). Graft ischemic time was typically 20–25 min and total operative time was 45–50 min with a success rate (beating hearts) of more than 90%. Rejection of heart grafts was not associated with death of recipients. The graft function was evaluated by palpation of the abdominal wall daily after operation. The function of the donor heart was assessed using a subjective score of 0 to 3 (zero for no beating; 0.5 for very weak beating; one for weak beating; two for moderate beating; three for full beating).

2.2.3 Plaque Assay

In this mouse model, virus titers in organs or sera were determined by Plaque Assay (Battegay et al., 1991; K. S. Lang, Recher, Navarini, et al., 2005). After harvesting and smashing organs in 1ml DMEM supplemented with 5% FCS, each cell suspension was transferred in 96-well plates and diluted with DMEM 5%FCS so that the cells were titrated at a ratio of 1:3. After transferring the samples in a

24- well plate, 200 μ l of freshly prepared MC57 cells diluted 8×10^5 cells/ml, were added. MC57 cells are susceptible for LCMV infection. Plates were ready for incubation and *Overlay* had to be added between 3 and 5 hours after starting incubation at 37°C and 5% CO₂. Incubation lasted for two days. At day three, cells were stained and evaluated for PFU. PFU were calculated according to the size of the smashed organ; dilution and logarithm of each variable is shown in the graph.

2.2.4 Flow Cytometry

Flow cytometry enables counting of fluorescent marked cells by 4 different lasers.

2.2.4.1 Tetramer Staining

Tetramers were provided by the National Institute of Health (NIH) Tetramer Facility. In short, 20 μ l of blood was stained with allophycocyanin (APC)-labeled GP33 MHC class I tetramers (GP33/H-2D^b) for 15 minutes at 37°C. After incubation, the samples were stained with anti-CD8 peridinin-chlorophyll-protein-complex (PerCP; BD Biosciences, Franklin Lakes, NJ) for 30 minutes at 4°C. Erythrocytes were then lysed using 1ml BD lysing solution (BD Biosciences); cells were washed once and analyzed by flow cytometry. Absolute numbers of GP33-specific CD8⁺ T cells per/ μ l blood were calculated by FACS analysis using fluorescent beads (BD Biosciences).

2.2.4.2 Intracellular Cytokine Staining (ICS)

For intracellular IFN- γ staining, splenocytes and lymphocytes from blood were stimulated with GP33 in the presence of Brefeldin A. After 6 hours, cells were stained for CD8 (eBioscience) for 30 minutes at 4°C, fixed with 2% Formaldehyde for 10 minutes and permeabilized with 1% Triton-X solution, and stained for IFN- γ with anti-mouse IFN- γ antibody (eBioscience) and analyzed by flow cytometry.

2.2.4.3 CFSE labeled Splenocyte Transfer

To follow proliferation of CD8⁺ TCR- specific T cells in blood, T cells were labeled with carboxyfluorescein succinimidyl ester (CFSE), a molecule, which made it possible to track cell proliferation due to decrease of tagged fluorescent molecule measured in FITC by flow cytometry. In this study, spleens of P14 mice have been smashed, suspended in PBS and filtered through a strainer. One spleen was suspended in 10 ml PBS. 5mM of CFSE has been added to the cell suspension and an incubation period of 10 minutes at 37°C followed. To eliminate remaining CFSE, single cell suspension has been washed twice with 10% FCS in DMEM. After discarding the supernatant, labeled cells were resuspended in IMDM to inject 200µl (10⁶ cells) intravenously in one mouse.

2.2.5 Depletion of CD8⁺ T Cells

For CD8⁺ T cell depletion, 500µg/100µl of anti-CD8 antibody clone YTS 169.4 (Bioxcell) was injected intraperitoneally on day -3 and -1 in bone marrow donor mice.

2.2.6 Triggering of Innate Immunity

Innate immune response was triggered by injecting 200µl of lipopolysaccharides (LPS) intraperitoneally (250µg/ml) in bone marrow chimera mice.

2.2.8 ELISA

Enzyme linked immunosorbent assay (ELISA) has been performed by the Mouse IL-10 ELISA Ready-SET-Go!® reagent kit using 20µl of plasma.

2.2.9 RT-PCR

Total RNA was isolated from organs with the TriFast®TRIzol (PeqLab, Erlangen) isolation method. Tissue samples with a weight range between 50-100mg have been homogenized in a TissueLyser (QIAGEN, Hilden). RNA was reverse-transcribed into cDNA with the QuantiTect Reverse Transcription Kit (QiAGEN, Hilden). Gene expression analysis was quantified by RT-PCR using SYBR®Green (AppliedBiosystems, Darmstadt) according to the protocol of the manufacturer. Expression levels were normalized against GAPDH and compared between study groups.

2.2.10 Analysis of Liver and Heart Enzymes

Serum and plasma were analysed in the central laboratory at the university hospital Essen for alanine transferase (ALT) and lactate dehydrogenase (LDH) and for CK-MB and Trop I. Serum or plasma has been collected from mice at various time points. 30µl serum or plasma has been diluted in 270µl PBS and were ready for measurement at the central laboratory.

3 Results

3.1 Transgenic LCMV-GP is ubiquitously expressed

To analyze immune responses of donor derived T cells in recipients expressing foreign antigen, *DEE* mice were used. *DEE* mice are transgenic mice which express LCMV-GP ubiquitously under the H2k promoter. The expression of LCMV-GP in the thymus, the organ of central tolerance induction was shown by RT-PCR (**Figure 3.1A**). LCMV-GP is a minor antigen which is presented on MHC class I. To confirm that the transgene LCMV-GP is presented by MHC class I molecules in host *DEE* mice, adoptive transfer of GP33 specific CD8⁺ T cells (P14 T cells) was performed. P14 cells showed proliferation in *DEE* mice (**Figure 3.1B**) but not in WT mice, when analyzed *ex vivo*.




Figure 3.1 LCMV-GP expression in *DEE* mice. A) RT-PCR analysis of LCMV-GP mRNA harvested from thymus of WT (n=2), *DEE* (n=3) and *Tcrb* deficient (LCMV; n=3) mice which were infected with 2×10^6 PFU of LCMV-WE representing positive control of LCMV-GP expression due to chronic infection at day 10 *post-infection*. Values show fold change to expression in WT mice. B) Proliferation of P14 T cells. Adoptive transfer of CFSE labeled P14 T cells into naïve WT mice and *DEE* mice were assessed by flow cytometry and shown in histogram blot (n=3).

To emphasize tolerance induction of LCMV-GP specific CD8⁺ T cells in mice carrying the minor antigen LCMV-GP, the F1 generation of P14 mice crossed with *DEE* mice were analyzed for GP33 specific CD8⁺ T cells. Due to negative selection, P14 T cells were deleted centrally (**Figure 3.2**).




Figure 3.2 Presence of GP33 specific CD8⁺ T cells in thymus and blood. A) Thymus of P14 mice (left panel) and P14x*DEE* mice (right panel) were gated on double positive CD4⁺/CD8⁺ T cells and for double positive CD90.2/Tet-GP33 T cells in the thymus. CD90.2 is expressed by mouse thymocytes and mature T cells. Results are shown as single dot plot. One dot plot is shown as a representative of each group (n=2). B) Number of CD8⁺ T cells in blood was determined in P14, P14x*DEE* and *DEE* mice shown in white columns and percentage of Tet-GP33 positive CD8⁺ T cells in black columns (n=4).

3.2 Host specific CD8⁺ T cells are not negatively selected after BMT

In this murine BMT model, P14 bone marrow was transferred in lethally irradiated *DEE* mice (**Figure 3.3**). P14 mice revealed LCMV-GP specific CD8⁺ T cells and *DEE* mice expressed LCMV-GP ubiquitously and present the antigen on MHC class I. LCMV-GP was used as a foreign antigen to demonstrate a single minor antigen mismatch.




Figure 3.3 Model of BMT. Recipient mice were lethally irradiated (9.5Gy) and received P14 bone marrow intravenously. *DEE* mice expressed GP33 ubiquitously. WT mice recipients were used as control. Mice were at least 6-8 weeks old.

As control, P14 bone marrow was transferred in WT mice. During reconstitution of bone marrow in recipients, blood was analyzed weekly for development of P14 T cells (**Figure 3.4A**). The graph shows the kinetic of GP33 specific CD8⁺ T cells in *DEE* and WT mice recipients after receiving P14 bone marrow. Even though, *DEE* recipients expressed GP33 everywhere in the tissue and present the antigen on MHC class I, P14 T cells remained ignorant. The mice did not become ill, indeed, measurement of cell damage marker such as LDH and ALT showed a physiological amount in sera (**Figure 3.4B**).




Figure 3.4 Development of P14 T cells after BMT and measurement of ALT and LDH in sera. A) Number of CD8⁺ T cells in blood was determined weekly in chimera mice by flow cytometry which is shown in the upper panel. According to the total number of CD8⁺ T cells, Tet-GP33 positive CD8⁺ T cells are shown in percentage in the panel below. B) ALT and LDH were determined in sera at indicated days post-transplant. Black squares represent mean values of B6 chimera mice at days post-transplant, which received P14 bone marrow *i.v.* (P14 BM→ WT) (n=4). Mean values of T cell number and ALT or LDH of irradiated *DEE* mice receiving P14 bone marrow (P14 BM→ *DEE*) are shown in white squares (n=4).

To characterize ignorant P14 T cells which developed in chimera mice more in detail, spleen has been removed after 40 days as reconstitution of bone marrow took approximately 30 days. Splenocytes were labeled with CFSE and transferred intravenously in different groups of mice. Splenocytes from WT and *DEE* chimera mice were injected *i.v.* in naïve WT mice, naïve *DEE* mice and WT mice infected with LCMV-WE 6 days *post-infection* to see whether P14 T cells behave differently in an environment of viral peptide presentation or active virus replication. P14 T cells which developed either in WT or *DEE* mice were transferred from spleen to WT, *DEE* or LCMV infected mice, proliferated in response to indigenous GP33-peptide (*DEE*) as well as to peptides from replicating virus (LCMV) (**Figure 3.5A**). *Ex vivo* restimulation of these cells with GP33-peptide to analyze intracellular cytokine staining (ICS) revealed no IFN- γ production in the absence of *in vivo*

antigen exposure (WT), low IFN- γ production after *in vivo* GP33-peptide exposure and significant IFN- γ production after *in vivo* exposure to replicating virus.

In conclusion P14 T cells proliferate in response to GP33-peptide presented in *DEE* mice but they are functionally ignorant. However, if these cells are exposed to an environment with actively replicating virus, these cells can be restimulated and produce IFN- γ .




Figure 3.5 P14 T cells generated in chimera mice got activated *ex-vivo*. A) 1×10^7 splenocytes of 40-days-old P14 BM→WT mice and P14 BM→DEE mice were labeled with CFSE and transferred into naïve WT mice, DEE mice and WT mice infected *i.v.* with 2×10^4 PFU of LCMV-WE 1 day after splenocyte transfer (n=2-4; pooled from two independent experiments). Proliferation of CD45.1⁺ CD8⁺ T cells (P14 T cells) derived from spleen was assessed by CFSE dilution 6 days after transfer. Histograms show cells gated on CD45.1⁺ CD8⁺ T cells in correlation to CFSE. One representative set of data is shown. B) Next, these splenocytes were analyzed for IFN- γ production after re-stimulation with the LCMV peptide GP33-41 *in vitro*. An intracellular cytokine staining was performed and CD45.1⁺ CD8⁺ specific T cells (P14) of either P14 BM→WT or P14 BM→DEE mice were characterized, IFN- γ secreting cells are given as percentage of lymphocytes (n=2-4; pooled from two independent experiments).

3.3 Introduction of LCMV-GP minor antigen in hematopoietic stem cells leads to deletion of host specific CD8⁺ T cells after BMT

During minor antigen mismatched BMT, host specific CD8⁺ T cells were present despite of minor antigen expression in the host. The hypothesis was that only antigen expressed in the hematopoietic derived cells is able to induce deletion of host specific CD8⁺ T cells in this setting. To test this hypothesis, we mixed P14 bone marrow cells with *DEE* bone marrow cells in a 1:1 ratio and transferred them into *DEE* and WT mice. In both groups, host specific CD8⁺ T cells were deleted (Figure 3.6)




Figure 3.6 P14 T cell kinetic in blood of chimeric mice after mixed bone marrow transplantation. Percentage of GP33 specific CD8⁺ T cells (P14 T cells) was determined in blood by flow cytometry at indicated time points after BMT (n=5).

3.4 Recipients ignore minor antigen (LCMV-GP) expressed in cardiac graft

Heterotopic HTX was performed as murine SOT model. Donor heart was harvested from *DEE* mice and transplanted in WT mice (**Figure 3.7**).




Figure 3.7 Model of heterotopic heart transplantation. GP33 expressing heart was harvested from *DEE* mice and transplanted in WT mice. As control, heart was harvested from WT mice and transplanted in WT mice.

Analyzing host versus graft reaction in SOT-setting showed that GP33 expression in the heart was completely ignored. Presence of minor antigen mismatch in syngeneic cardiac graft did not induce rejection in a time period of 100 days whereas allogeneic hearts harvested from Sv/129 mice were rejected in WT mice within 10 days (**Figure 3.8**). Sv/129 mice differ genetically from WT mice by MHC (H2 complex) class I and II alleles and the genetic background and were used as allograft rejection model (positive control).




Figure 3.8 Graft survival curve after HTX. Graft survival was defined as persistence of heart beating. Graft survival was independent of survival of recipient mice. B6 (black square), *DEE* (GP33; white square) and Sv/129 (black triangle) hearts were harvested as donor hearts and transplanted in B6 (WT) mice. Heart beat was determined weekly by abdominal palpation (n=4-5).

Moreover, no development of GP33 specific CD8⁺ T cells could be detected in WT mice recipients transplanted with a syngeneic WT heart or a *DEE* heart carrying the LCMV-GP33 peptide (**Figure 3.9**). No priming of CD8⁺ T cells took place. In conclusion, minor antigen mismatch was completely ignored by host derived WT CD8⁺ T cells, in this setting.




Figure 3.9 Tetramer staining of blood after HTX. At day 100 post-transplant blood from transplanted mice receiving either B6 (WT) heart or *DEE* (GP33) heart were analyzed for GP33 specific CD8⁺ T cells by flow cytometry. Dotplot was gated on lymphocytes. Left panel shows representative dotplot of B6 (WT) mice receiving B6 (WT) heart (n=3). Right panel shows representative dotplot of B6 (WT) mice receiving *DEE* heart (n=4).

3.5 Immunological tolerance in an LCMV model

WT mice received either naïve B6 heart or LCMV loaded heart. Effect of viral transmitted infection by HTX versus systemic (*i.v.*) LCMV infection post-transplantation has been analyzed as followed (**Figure 3.10**).




Figure 3.10 Experiment design. WT mice recipients received either a naïve heart transplant or LCMV loaded heart. WT mice receiving a naïve heart transplant were infected with LCMV strain WE intravenously.

3.5.1 LCMV carrier mice

To create LCMV carrier mice, *Tcrb*^{-/-} mice which cannot mount an antiviral immune response because of TCR deficiency required for MHC class I expression, have been infected with 2×10^6 PFU of LCMV strain WE. Mice were then chronically infected and virus persisted in mice. Virus titer is shown and is comparable to LCMV-WE high dose infection (**Figure 3.11**). LCMV carrier heart was used as donor heart which was severely infected with replicating virus. In the previous experiments (**Figure 3.5**) GP33 expression alone was not able to induce immune activation. The hypothesis was that only replicating virus was able to break immunological ignorance.




Figure 3.11 Virus titers in LCMV donor hearts. *Tcrb*^{-/-} mice were infected with 2×10^6 PFU of LCMV strain WE. Graph shows viral titers in heart after 30 days post-infection (n = 4). Horizontal dotted lines designate the detection limit.

3.5.2 LCMV transmitted by HTX leads to T cell exhaustion in WT mice

In this setting, viral loaded hearts were transplanted in naïve mice and naïve hearts were transplanted in virus infected mice. The syngeneic grafts whether naïve or infected were not rejected. The hearts kept on beating and recipients survived. WT mice receiving chronically infected hearts became carrier mice; meaning, virus persisted in mice and spread from the transplanted heart to other tissue. Mice receiving naïve-HTX and were infected systemically with LCMV cleared the virus. Plaque assay of organs and heart transplant showed viral

persistence 30 days post-transplant (**Figure 3.12**) in LCMV-HTX mice but not in LCMV infected naïve-HTX mice.




Figure 3.12 Virus titers in organs after LCMV-HTX. Organs have been removed 30 days post-transplant for Plaque Assay. White columns show virus titers in spleen, liver, kidney, lung, transplanted heart (Tx) and endogenous heart (Endo) removed from WT mice recipients receiving LCMV-loaded hearts (n=4). Control group (black columns) received naïve hearts and were infected with LCMV strain WE intravenously 15 days post-transplant. 30 days post-infection organs have been harvested for Plaque Assay (n=4). Horizontal dotted lines designate the detection limit.

Virus infection could not be controlled in LCMV-HTX mice. Analyzing virus specific CD8⁺ T cells, which normally develop after LCMV-WE infection, revealed a reduced GP33 specific CD8⁺ T cell number in blood (**Figure 3.13A**). Furthermore viral specific response in form of effector cytokine production such as IFN- γ was reduced in blood (**Figure 3.13B**). Together these data demonstrate anti-viral T cell exhaustion in chronically infected mice. T cell exhaustion is defined as a loss of effector function (Wherry, 2011; Yi et al., 2010). Gradual loss can be distinguished as stages which are determined by diminished or vanished production of effector cytokines such as IFN- γ , increased expression of inhibitory receptors and deletion of antigen-specific T cells. In addition, T cell exhaustion correlates with viral persistence.




Figure 3.13 CD8⁺ T cell kinetic and IFN-γ production in blood. A) Percentage of GP33 specific CD8⁺ T cells was determined in blood by flow cytometry at indicated time points post-transplant or post-infection respectively. WT mice received LCMV carrier hearts (white squares; n=4). Control group consisted of WT mice which received naïve hearts and were infected *i.v.* with LCMV strain WE low dose 15 days post-transplant (black squares; n=4). Time is shown in days post-transplant and post-infection respectively. B) IFN-γ production was assessed by intracellular cytokine staining (ICS) in blood gated on CD8⁺ T cells. White columns represent WT mice receiving LCMV loaded hearts (n=4) at days post-transplant and as control, naïve WT mice infected with LCMV strain WE high dose at days 8 and 14 post-infection.

3.6 Breaking immunological tolerance

In the study reported here, minor antigen expression in hosts (*DEE* mice) receiving BMT, induced ignorance in donor BM derived minor antigen specific CD8⁺ T cells (P14 cells). Antigen expression in syngeneic graft (*DEE* heart) during SOT was ignored by the lymphoid system of the host (WT mice). Replicating antigen in transplant (LCMV- HTX) induced tolerance in naïve transplant recipients due to T cell exhaustion. Now the question occurred how to break this tolerance.

3.6.1 High frequencies of host specific CD8⁺ T cells induce limited GvHD after stimulation with LPS

Donor derived LCMV-GP minor antigen specific CD8⁺ T cells (P14) grafted in *DEE* mice remained ignorant, which means, these mice did not develop GvHD. However, innate immune activation via Toll-like Receptor (TLR) can have an impact on adaptive auto-reactive immune responses (K. S. Lang et al., 2006; K. S. Lang, Recher, Junt, et al., 2005).

Therefore, chimera mice which received P14 bone marrow were treated with lipopolysaccharides (LPS). LPS was injected intraperitoneally in P14 BM→WT mice and P14 BM→*DEE* mice 200 days after BMT. Serologically, LDH was measured as general marker for cell damage and transaminases as specific markers for liver disease. When LPS was administrated LDH was enhanced in P14 BM→*DEE* mice but not in P14 BM→WT mice, indicating cell damage took place (**Figure 3.14**). Specific liver enzymes were in normal range. This suggests that LPS impacts on immune activation of host specific CD8⁺ T cells. However, these data show that even at high precursor frequencies of minor antigen specific CD8⁺ T cells acute GvHD is not induced.




Figure 3.14 ALT and LDH measurement in serum after LPS challenge. P14 bone marrow was transferred in *DEE* and WT mice (white columns P14 BM in *DEE* mice; black columns P14 BM in WT mice (n=4-5) pooled from two experiments). 50µg LPS was injected intraperitoneally 200 days after BMT. 1 and 2 days after LPS challenge, serum was analyzed for ALT and LDH amount.

3.6.2 IL-10 deficient mice die after transplantation of LCMV infected hearts

Generation of GP33 specific CD8⁺ T cells in the presence of *DEE* transplanted heart did not take place due to T cell exhaustion (**Figure 3.8**). Possible mechanisms of CD8⁺ T cell exhaustion during heart transplantation can vary. In this study, CD8⁺ T cell exhaustion led to tolerance of heart transplant and consequently to syngeneic graft survival. Interleukin-10 is known to play a crucial role for CD8⁺ T cell exhaustion during chronic infection (Blackburn et al., 2007; Brooks et al., 2006). Therefore, IL-10 deficient mice were used as recipient (**Figure 3.15**).




Figure 3.15 Experiment design. LCMV carrier hearts were transplanted in *IL-10*^{-/-} mice recipients or in WT mice recipients. Time line shows the days post-transplant. At indicated time points, blood analysis was done by flow cytometry.

Interleukin-10 is an immunosuppressive cytokine which is secreted by various types of cells such as CD8⁺ and CD4⁺ T cells, B cells and APCs (Moore et al., 2001). When mice lack IL-10 during LCMV-HTX, they died (**Figure 3.16A**). This can be due to high numbers of activated CD8⁺ T cells. Thus, infection with LCMV-WE intravenously in IL-10 deficient mice led to fast generation of LCMV specific CD8⁺ T cells (data not shown). When LCMV carrier hearts were transplanted into IL-10 deficient mice, higher numbers of CD8⁺ T cells could be detected in blood (**Figure 3.16B**). Indeed, IL-10 was elevated in WT mice receiving an infected heart transplant (**Figure 3.17A**). Virus titer in blood was not affected by IL-10 loss (**Figure 3.17B**).




Figure 3.16 Survival curve and CD8⁺ T cell kinetic after LCMV-HTX. LCMV carrier hearts were transplanted in *IL-10*^{-/-} mice recipients (white circle) and in WT mice recipients (black square). A) Mice were observed in a time period of 30 days. Survival of mice recipients after LCMV-HTX is shown in percentage (n=7-15; pooled from 2 to 3 experiments). B) CD8⁺ T cells were measured in blood by flow cytometry at indicated time points post-transplant (n=2-4).




Figure 3.17 IL-10 and virus titers in plasma and blood after LCMV-HTX. LCMV carrier hearts were transplanted in *IL-10*^{-/-} mice recipients and in WT mice recipients (n=2-4). A) IL-10 was measured in plasma by ELISA at day 2 and 6 post-transplant. B) Plaque Assay was performed from blood. Virus load was determined by plaque-forming units in blood at days post-transplant. Horizontal dotted lines designate the detection limit.

IL-10 deficient mice rejected the hearts within 10 days and heart enzymes were elevated in *IL-10*^{-/-} mice (**Figure 3.18A**). In addition, *IL-10*^{-/-} mice became ill and revealed high levels of LDH and ALT indicating massive cell death in tissue and liver damage respectively (**Figure 3.18B**). The mice revealed a systemic illness and died within 10 days due to immunopathology. Together these data show that

IL-10 was essential to prevent syngeneic graft rejection after transplantation of viral infected hearts.




Figure 3.18 Analysis of heart and liver enzymes in sera after LCMV-HTX. LCMV carrier hearts were transplanted in *IL-10*^{-/-} mice recipients and in WT mice recipients. A) CK-MB and Trop I (specific heart enzymes) were measured in sera at day 4, 6 and 8 post-transplant (n=2-4). B) ALT and LDH (marker for liver damage and cell death) were determined in sera at day 4, 6 and 8 post-transplant (n=2-4).

3.7 Graft rejection by memory T cells

Are memory T cells able to prevent virus persistence? Are they dependent on IL-10? Therefore, memory CD8⁺ T cells were generated in C57BL/6 mice by infecting mice with LCMV-WE intravenously. After priming with LCMV, mice generated a specific immune response. In a next step, LCMV carrier hearts were transplanted into LCMV infected mice (**Figure 3.19**).




Figure 3.19 Experiment design. Naïve C57BL/6 (WT) mice were infected at least 20 days before HTX with 200 PFU of LCMV-WE intravenously and generated memory CD8⁺ T cells. Memory mice or naïve C57BL/6 (WT) mice received LCMV infected hearts and blood analysis were made at indicated time points post-transplant.

3.7.1 Memory mice recipients reject LCMV infected heart transplant

Transplantation of LCMV infected hearts led to massive CD8⁺ T cell expansion in memory mice (**Figure 3.20A**). Memory CD8⁺ T cells got highly activated as shown by down regulation of IL-7 receptor (CD127) and CD62L (**Figure 3.20B**). Virus specific memory CD8⁺ T cells could control LCMV (**Figure 3.21A**) and transplanted hearts were early rejected in memory mice (**Figure 3.21B**). In line with the acute heart rejection, heart specific parameters, CK-MB and Trop I, raised early after transplantation (**Figure 3.21C**). Histology revealed high numbers of infiltrating CD8⁺ T cells in the heart (data not shown). In conclusion, we found that presence of memory CD8⁺ T cells resulted in fast rejection of viral infected hearts, but virus could be controlled and the mice survived.




Figure 3.20 Characterization of Memory T cells. A) Graph showing frequency of T cells that were positive for the MHC class I tetramer of the glycoprotein of LCMV (Tet-GP33⁺) and for CD8 (CD8⁺) in blood of WT or LCMV infected recipients receiving LCMV loaded hearts shown in days post-transplant (n=3-5). B) Analysis of activation marker such as CD127, CD62L, CD44 and PD-1 are made by gating TET-GP33⁺ CD8⁺ T cells and shown in a histogram blot at day 6 post-transplant in comparison to day -1 pre-transplant in blood. One histogram blot is shown as a representative of each group.




Figure 3.21 Virus control and transplant rejection in memory mice. LCMV carrier hearts were transplanted in memory mice recipients and in WT mice. A) Virus titers in blood were determined by plaque-forming units (PFU) in Plaque assay at days 7 and 16 post-transplant. Horizontal dotted lines designate the detection limit (n=4). B) Allograft survival curve after LCMV-HTX was determined by rest of heart beat (n=4-7). Allograft survival was independent of recipient survival. C) Measurement of heart enzymes (CK-MB and Trop I) were determined in sera (n=3-4).

4 Discussion

In the study reported here, the role of viral infection in transplantation medicine was assessed in a murine LCMV model. Two main experimental but clinical relevant conditions have been used: murine bone marrow transplantation and heterotopic heart transplantation. Minor antigens as well as viral infection influence outcome of patients undergoing BMT or SOT in fully matched donor and recipients. The LCMV-GP transgene was used as a peptide in a minor antigen mismatch transplantation model. In a next step, the role of replicating antigen was analyzed in heterotopic heart transplantation. LCMV transmitted by heart transplantation led to T cell exhaustion in WT mice and in consequence to syngeneic graft tolerance. Moreover, induction of host antigen (GP33) in hematopoietic stem cells led to deletion of host specific CD8⁺ T cells after BMT.

4.1 Host specific CD8⁺ T cells proliferate in presence of minor antigen in host after BMT

In the murine minor antigen mismatched bone marrow model we transplanted P14 bone marrow into *DEE* mice to analyze the activation of these host specific CD8⁺ T cells. The recipients developed high frequencies of GP33-specific T-lymphocytes independent of antigen expression in the thymus and peripheral organs. Nevertheless, they did not develop lethal disease in form of GvHD. To confirm central tolerance induction by recognition of GP33 minor antigen and by T cell deletion we crossed P14 mice with *DEE* mice. Almost no GP33-specific CD8⁺ T cell has been detected in peripheral organs. *DEE* mice express ubiquitously GP33 peptide in organs and transferring CFSE labeled mature but naïve P14 T cells into *DEE* mice led to significant T cell proliferation, suggesting that the minor antigen is detectable by the transgenic TCR of the P14 T cells. As described in previous publications proliferation of anti-host reactive CD8⁺ T cells could be induced by minor antigen presentation in the host (Mori et al., 2005). It

remains unclear, whether presentation of minor antigens in the host can lead to ignorance and finally to reduced immunopathology in patients after BMT.

4.2 Triggering the innate immune system can lead to activation of host specific CD8⁺ T cells

We demonstrated that triggering the innate immune system with LPS could lead to significant cell damage in *DEE* recipients whereas WT recipients remained unaffected by GP33-specific CD8⁺ T cells. This is in line with the clinical situation where bacterial or viral infection (e.g. CMV) often induces GvHD (Ljungman, 2008). Especially for the human cytomegalovirus (HCMV), preexisting data show an association of viral infection with increased risk of aGvHD (Boeckh et al., 2004; Nichols et al., 2002). Interestingly, in a clinical trial it was shown that early replicative HCMV infection reduces the risk of leukemic relapse in patients with acute myeloid lymphoma (AML) (Elmaagacli et al., 2011). Though the underlying mechanism is not completely understood yet; a virus-induced abrogation of immune ignorance comparable to our murine model and even induction of graft versus leukemia (GvL) is a possible explanation for reduced risk of leukemic relapse after early replicative HCMV infection. Thus, beyond genetic haplotype matches the immune status on bacterial and viral infections of bone marrow donors might be helpful for GvHD risk evaluation, which is already performed for herpes viruses and HCMV (Pietersma et al., 2011).

4.3 Outlook for possible therapeutic methods in BMT

These data are in contrast to current models, suggesting that low frequencies of anti-host reactive CD8⁺ T cells result in development of GvHD (Desmarests et al., 2009; Schroeder et al., 2011). We found that in absence of secondary stimulation even high numbers of host-reactive T cells are ignorant. GvHD is often associated with GvL. GvL can result in complete control of malignant

cells after minor HLA mismatch. Interestingly efficient GvL can occur even in the presence of mild GvHD (Klymenko et al., 2011). The reasons for that are not well explained. From our data we would suggest that the presentation of leukemic antigens in primary lymphoid organs is crucial for the recognition by anti-host reactive CD8⁺ T cells and for early elimination. Therefore, a promising approach to treat GvHD would be the inhibition of T cell egress to peripheral organs. This would potentially limit aGvHD, but still allow GvL. Such a preparation is Fingolimod (FTY720) which is licensed for treatment of relapsing remitting multiple sclerosis. Fingolimod prevents the egress of lymphocytes from secondary lymphoid organs thereby preventing migration of immune cells to the central nervous system (Chun et al., 2010). Fingolimod is not clinically tested in BMT. Even though its application seems attractive, Fingolimod has first to be evaluated in animal models since it reduces the peripheral lymphocyte count and acts as an immunosuppressant. In conclusion we demonstrated that high frequencies of anti-host reactive CD8⁺ T cells can remain ignorant after engraftment in recipient and do not induce GvHD.

4.4 Immunized recipients have an increased risk for cardiac graft rejection

We mimicked the clinical situation of donor-derived viral infection in SOT in form of heterotopic heart transplantation in a mouse model. This system allows us to focus on recipients' immune response against virus infection transmitted through organ transplantation.

Clinically, CMV is a major problem in SOT (Razonable et al., 2013). CMV is a herpesvirus which is widely spread in human beings. Seroprevalence of CMV varies between 30 and 97% in the population (Bate et al., 2010; Cannon et al., 2010). Consequently, more than 50% of solid organ transplant recipients suffer from CMV infection in the first 3 months post-transplant. Seropositivity of donor and recipient play a crucial role in morbidity and mortality for the transplant recipient. CMV-seropositive donors and recipients are designated as D⁺ and R⁺

respectively. According to seroprevalence, different risk groups are categorized. In the D⁺ and R⁻ (D⁺/R⁻) group the recipient has the highest risk for CMV disease. The lowest risk for CMV disease is in the case of seronegativity in donor and recipient (D⁻/R⁻) (Ramanan et al., 2013). A CMV prophylaxis treatment is preserved for transplant recipients who are at highest risk for CMV disease (D⁺/R⁻) and (D⁺/R⁺). A retrospective study compared morbidity and mortality in four CMV donor/recipient serostatus categories (Harvala et al., 2013). As result, the group D⁺/R⁺ showed the highest risk for mortality post-transplant, even though CMV disease occurs more often in D⁺/R⁻. In line with our data, recipients who have been infected once before transplantation (R⁺) show a strong immune response against the allograft (D⁻) leading to rejection. In our study, LCMV infection via transplant (D⁺) in naïve recipient (R⁻) mice led to chronic virus infection due to CD8⁺ T cell exhaustion. The grafts survived and showed only partial dysfunction.

4.5 Role of LCMV infection in human beings

Another example for donor-derived infection in humans is LCMV. LCMV is an arenavirus, which is very rare in human beings. An infection is usually asymptomatic in healthy persons and can be caused after close contact to house mice or hamsters (Childs et al., 1992; Childs et al., 1991). Some cases are known where LCMV was transmitted during transplantation. Donor-derived infection in several recipients led to death within three weeks after transplantation. The same viral strain could be detected in a cluster of patients, where seven of eight recipients died. The recipients revealed unspecific symptoms such as diarrhea, fever and systemic illness (Fischer et al., 2006). Normally, LCMV infection in non-immunosuppressed people is asymptomatic.

Immunosuppression after transplantation aims to reduce immune response by the adaptive immune system. Nevertheless, it encloses many side effects such as a higher risk of opportunistic infections, a heightened incidence of cancer and toxicity (Halloran, 2004). In our study, viral infected hearts were transplanted in

immunocompetent mice and were fully HLA-matched to focus specifically on CD8⁺ immune response.

4.6 T cell exhaustion- a possible mechanism inducing tolerance?

A new perspective for tolerance in SOT is to understand CD8⁺ T cell exhaustion. Little is known in the field of transplantation medicine. We know that T cell exhaustion limits the immune response against chronic infections and tumors. In case of recrudescence of exhausted T cells, clearance of infections and tumor regression can be induced (Blank et al., 2007).

Previous clinical studies effort to reactivate exhausted T cells as treatment for chronic infections or tumor disease (Sharpe et al., 2015). However, the mechanisms leading to T-cell exhaustion are unknown. Further, the role of T-cell exhaustion in transplantation or autoimmunity is not defined. Recently, *Sarraj et al* (Sarraj et al., 2014) showed that genetic deletion of selectin ligand for leucocyte migration leads to impaired T cell function. They focused on CD4⁺ T cells and induced T cell exhaustion in a murine transplantation model by impaired migration capability. They determined T cell exhaustion by impaired effector cytokine production (IFN- γ), defective proliferation of CD4⁺ T cells and higher PD-1 expression. Knock-out mice recipients providing impaired migration capability revealed T cell exhaustion and showed prolonged graft survival. T cells were not able to migrate in the MHC II mismatched heart transplant which in consequence led to permanent activation of T cells and then to T cell exhaustion. Graft survival was significantly increased. The data demonstrate that T cell exhaustion led to tolerance of allograft.

The fact that IL-10 is an immunoregulatory cytokine that is associated with T cell exhaustion, is in line with the above finding (Blackburn et al., 2007). Its function during virus infection is described to be immunosuppressive by limiting cytokine production and proliferation of CD8⁺ and CD4⁺ T cells leading to viral persistence. A variety of cells can produce IL-10 such as T cells, B cells and APCs (Blackburn et al., 2007; Moore et al., 2001). Clinical studies demonstrated that

polymorphisms linked with increased IL-10 production are associated with increased susceptibility to chronic HCV infection and increased severity of chronic HBV infection (Knapp et al., 2003; Paladino et al., 2006; Persico et al., 2006). Vice versa, polymorphisms with reduced expression of IL-10 correlate with a slower progression of AIDS in HIV- infected patients (Shin et al., 2000). Its role in transplantation medicine is poorly understood. As IL-10 is known for its anti-inflammatory response and immunosuppressive role, it should abet graft survival. Indeed, IL-10 inhibits ischemia/reperfusion injury (Deng et al., 2001), extends allograft survival and function (Cypel et al., 2009; Feng et al., 1999; Zuo et al., 2001) and is essential for the action of regulatory T cells mediating tolerance at least in some transplant models (Hara et al., 2001). In our model, IL-10 deficient mice receiving LCMV loaded heart died early after transplantation. Loss of IL-10 in recipients showed a systemic immune response with immunopathology indicating that absence of IL-10 prevents T cell exhaustion. Its immunoregulatory function in virus infection through organ transplantation helps tolerating the allograft.

This changed in case of memorized mice. Memory CD8⁺ T cells were capable to control donor-derived LCMV infection but induced an acute graft rejection. Memory T cells play a major role in acute and chronic allograft rejection. Pre-transplant frequency of donor-specific alloreactive memory T cells in recipient correlates with the risk of long-term allograft rejection (Heeger et al., 1999). Previous experimental models demonstrate their potential alloreactivity. They even showed that cross-reactivity can lead to alloreaction defined as heterologous immunity (van den Heuvel et al., 2015). In our case replication competent LCMV in heart transplant was rejected in memory mice. These facts emphasize the significance of donor-derived viral infection. Immunized patients exposed to viral load facing possible alloreaction have to be treated virus specifically (e.g. CMV). It decreases significantly graft survival in SOT. IL-10 therapy in case of unexpected donor-derived viral infection could be a potential immunotherapy in SOT.

5 Summary

Graft versus host disease (GvHD) occurs in 40% of cases with patients having a MHC I matched bone marrow transplantation (BMT). Mechanisms causing this disease remain to be studied. Here we used a CD8⁺ T cell transgenic mouse strain (P14/CD45.1⁺) and *DEE* mice bearing the foreign antigen (LCMV-GP33-41) to study mechanisms of tolerance in donor derived host specific CD8⁺ T cells after BMT. We found that host reactive CD8⁺ T cells were not negatively selected in the thymus and developed comparably to host non-specific CD8⁺ T cells. Host specific CD8⁺ T cells ignored the antigen expressed ubiquitously by host cells but they could be activated *ex vivo* via LCMV-infection. Lipopolysaccharides (LPS) induced transient cell damage in *DEE* mice bearing host specific CD8⁺ T cells, suggesting that induction of host inflammatory response could break this ignorance. In conclusion, we found that after BMT host specific CD8⁺ T cells ignore antigen in recipients and that they are only deleted when host antigen is present in the hematopoietic system. Moreover, LPS-induced immune activation contributes to induction of alloreactivity of host specific CD8⁺ T cells after BMT. Unexpected transmissions of viral pathogens during solid organ transplantation (SOT) can result in severe, life-threatening diseases in transplant recipients. Immune activation contributes to disease onset; however mechanisms balancing the immune response against transmitted virus infection through organ transplantation remain unknown. Here, we found, using LCMV, that transplantation of LCMV infected hearts led to exhaustion of virus specific CD8⁺ T cells, viral persistence in organs and survival of graft and recipient. Genetic depletion of IL-10 resulted in a strong immune activation, graft dysfunction and death of mice, suggesting that IL-10 was a major regulator of CD8⁺ T cell exhaustion during SOT. In the presence of memory CD8⁺ T cells, virus could be controlled; however sufficient antiviral immune response resulted in rejection of transplanted heart. In conclusion, we found that virus transmitted by SOT cannot be controlled by naive recipients due to IL-10 mediated CD8⁺ T cell exhaustion which thereby prevented immunopathology and graft failure whereas memory mice recipients were able to control the virus and induced graft failure.

6 References

1. Abdul Wahid, S. F., Ismail, N. A., Mohd-Idris, M. R., Jamaluddin, F. W., Tumian, N., Sze-Wei, E. Y., Muhammad, N., Nai, M. L. (2014). Comparison of reduced-intensity and myeloablative conditioning regimens for allogeneic hematopoietic stem cell transplantation in patients with acute myeloid leukemia and acute lymphoblastic leukemia: a meta-analysis. *Stem Cells Dev* 23, 2535-2552.
2. Ahmed, R., Oldstone, M. B. (1988). Organ-specific selection of viral variants during chronic infection. *J Exp Med* 167, 1719-1724.
3. Bate, S. L., Dollard, S. C., Cannon, M. J. (2010). Cytomegalovirus seroprevalence in the United States: the national health and nutrition examination surveys, 1988-2004. *Clin Infect Dis* 50, 1439-1447.
4. Battegay, M., Cooper, S., Althage, A., Banziger, J., Hengartner, H., Zinkernagel, R. M. (1991). Quantification of lymphocytic choriomeningitis virus with an immunological focus assay in 24- or 96-well plates. *J Virol Methods* 33, 191-198.
5. Blackburn, S. D., Wherry, E. J. (2007). IL-10, T cell exhaustion and viral persistence. *Trends Microbiol* 15, 143-146.
6. Blank, C., Mackensen, A. (2007). Contribution of the PD-L1/PD-1 pathway to T-cell exhaustion: an update on implications for chronic infections and tumor evasion. *Cancer Immunol Immunother* 56, 739-745.
7. Boeckh, M., Nichols, W. G. (2004). The impact of cytomegalovirus serostatus of donor and recipient before hematopoietic stem cell transplantation in the era of antiviral prophylaxis and preemptive therapy. *Blood* 103, 2003-2008.
8. Borst, J., Jacobs, H., Brouns, G. (1996). Composition and function of T-cell receptor and B-cell receptor complexes on precursor lymphocytes. *Curr Opin Immunol* 8, 181-190.
9. Brooks, D. G., Trifilo, M. J., Edelmann, K. H., Teyton, L., McGavern, D. B., Oldstone, M. B. (2006). Interleukin-10 determines viral clearance or persistence in vivo. *Nat Med* 12, 1301-1309.
10. Call, M. E., Wucherpfennig, K. W. (2005). The T cell receptor: critical role of the membrane environment in receptor assembly and function. *Annu Rev Immunol* 23, 101-125.
11. Cannon, M. J., Schmid, D. S., Hyde, T. B. (2010). Review of cytomegalovirus seroprevalence and demographic characteristics associated with infection. *Rev Med Virol* 20, 202-213.

12. Champlin, R. E., Passweg, J. R., Zhang, M. J., Rowlings, P. A., Pelz, C. J., Atkinson, K. A., Barrett, A. J., Cahn, J. Y., Drobyski, W. R., Gale, R. P., Goldman, J. M., Gratwohl, A., Gordon-Smith, E. C., Henslee-Downey, P. J., Herzig, R. H., Klein, J. P., Marmont, A. M., O'Reilly, R. J., Ringden, O., Slavin, S., Sobocinski, K. A., Speck, B., Weiner, R. S., Horowitz, M. M. (2000). T-cell depletion of bone marrow transplants for leukemia from donors other than HLA-identical siblings: advantage of T-cell antibodies with narrow specificities. *Blood* 95, 3996-4003.
13. Childs, J. E., Glass, G. E., Korch, G. W., Ksiazek, T. G., Leduc, J. W. (1992). Lymphocytic choriomeningitis virus infection and house mouse (*Mus musculus*) distribution in urban Baltimore. *Am J Trop Med Hyg* 47, 27-34.
14. Childs, J. E., Glass, G. E., Ksiazek, T. G., Rossi, C. A., Oro, J. G., Leduc, J. W. (1991). Human-rodent contact and infection with lymphocytic choriomeningitis and Seoul viruses in an inner-city population. *Am J Trop Med Hyg* 44, 117-121.
15. Chun, J., Hartung, H. P. (2010). Mechanism of action of oral fingolimod (FTY720) in multiple sclerosis. *Clin Neuropharmacol* 33, 91-101.
16. Complete sequence and gene map of a human major histocompatibility complex. The MHC sequencing consortium. (1999). *Nature* 401, 921-923.
17. Corry, R. J., Winn, H. J., Russell, P. S. (1973). Heart transplantation in congenic strains of mice. *Transplant Proc* 5, 733-735.
18. Cypel, M., Liu, M., Rubacha, M., Yeung, J. C., Hirayama, S., Anraku, M., Sato, M., Medin, J., Davidson, B. L., de Perrot, M., Waddell, T. K., Slutsky, A. S., Keshavjee, S. (2009). Functional repair of human donor lungs by IL-10 gene therapy. *Sci Transl Med* 1, 4ra9.
19. Davis, C. L., Hricik, D. E. (2004). Transplant: immunology and treatment of rejection. *Am J Kidney Dis* 43, 1116-1137.
20. Dempsey, P. W., Vaidya, S. A., Cheng, G. (2003). The art of war: Innate and adaptive immune responses. *Cell Mol Life Sci* 60, 2604-2621.
21. Deng, J., Kohda, Y., Chiao, H., Wang, Y., Hu, X., Hewitt, S. M., Miyaji, T., McLeroy, P., Nibhanupudy, B., Li, S., Star, R. A. (2001). Interleukin-10 inhibits ischemic and cisplatin-induced acute renal injury. *Kidney Int* 60, 2118-2128.
22. Dengler, T. J., Zimmermann, R., Braun, K., Muller-Bardorff, M., Zehelein, J., Sack, F. U., Schnabel, P. A., Kubler, W., Katus, H. A. (1998). Elevated serum concentrations of cardiac troponin T in acute allograft rejection after human heart transplantation. *J Am Coll Cardiol* 32, 405-412.

23. Desmarests, M., Cadwell, C. M., Peterson, K. R., Neades, R., Zimring, J. C. (2009). Minor histocompatibility antigens on transfused leukoreduced units of red blood cells induce bone marrow transplant rejection in a mouse model. *Blood* 114, 2315-2322.
24. Dong, V. M., Womer, K. L., Sayegh, M. H. (1999). Transplantation tolerance: the concept and its applicability. *Pediatr Transplant* 3, 181-192.
25. Elmaagacli, A. H., Steckel, N. K., Koldehoff, M., Hegerfeldt, Y., Trenchel, R., Ditschkowski, M., Christoph, S., Gromke, T., Kordelas, L., Ottinger, H. D., Ross, R. S., Horn, P. A., Schnittger, S., Beelen, D. W. (2011). Early human cytomegalovirus replication after transplantation is associated with a decreased relapse risk: evidence for a putative virus-versus-leukemia effect in acute myeloid leukemia patients. *Blood* 118, 1402-1412.
26. Feng, X., Zheng, X. X., Yi, S., Lehnert, A. M., Strom, T. B., O'Connell, P. J. (1999). IL-10/Fc inhibits macrophage function and prolongs pancreatic islet xenograft survival. *Transplantation* 68, 1775-1783.
27. Ferrara, J. L., Deeg, H. J. (1991). Graft-versus-host disease. *N Engl J Med* 324, 667-674.
28. Ferrara, J. L., Levine, J. E., Reddy, P., Holler, E. (2009). Graft-versus-host disease. *Lancet* 373, 1550-1561.
29. Fischer, S. A., Graham, M. B., Kuehnert, M. J., Kotton, C. N., Srinivasan, A., Marty, F. M., Comer, J. A., Guarner, J., Paddock, C. D., DeMeo, D. L., Shieh, W. J., Erickson, B. R., Bandy, U., DeMaria, A., Jr., Davis, J. P., Delmonico, F. L., Pavlin, B., Likos, A., Vincent, M. J., Sealy, T. K., Goldsmith, C. S., Jernigan, D. B., Rollin, P. E., Packard, M. M., Patel, M., Rowland, C., Helfand, R. F., Nichol, S. T., Fishman, J. A., Ksiazek, T., Zaki, S. R., Team, L. i. T. R. I. (2006). Transmission of lymphocytic choriomeningitis virus by organ transplantation. *N Engl J Med* 354, 2235-2249.
30. Fishman, J. A. (2007). Infection in solid-organ transplant recipients. *N Engl J Med* 357, 2601-2614.
31. Fishman, J. A., Strong, D. M., Kuehnert, M. J. (2009). Organ and tissue safety workshop 2007: advances and challenges. *Cell Tissue Bank* 10, 271-280.
32. Fujii, M., Hara, H., Meng, W., Vonsattel, J. P., Huang, Z., Moskowitz, M. A. (1997). Strain-related differences in susceptibility to transient forebrain ischemia in SV-129 and C57black/6 mice. *Stroke* 28, 1805-1810; discussion 1811.

33. Funabashi, H., Uchida, K., Kado, S., Matsuoka, Y., Ohwaki, M. (2001). Establishment of a Tcrb and Trp53 genes deficient mouse strain as an animal model for spontaneous colorectal cancer. *Exp Anim* 50, 41-47.
34. Gerlai, R. (1996). Gene-targeting studies of mammalian behavior: is it the mutation or the background genotype? *Trends Neurosci* 19, 177-181.
35. Grossi, P. A., Fishman, J. A., Practice, A. S. T. I. D. C. o. (2009). Donor-derived infections in solid organ transplant recipients. *Am J Transplant* 9 Suppl 4, S19-26.
36. Halloran, P. F. (2004). Immunosuppressive drugs for kidney transplantation. *N Engl J Med* 351, 2715-2729.
37. Hara, M., Kingsley, C. I., Niimi, M., Read, S., Turvey, S. E., Bushell, A. R., Morris, P. J., Powrie, F., Wood, K. J. (2001). IL-10 is required for regulatory T cells to mediate tolerance to alloantigens in vivo. *J Immunol* 166, 3789-3796.
38. Harvala, H., Stewart, C., Muller, K., Burns, S., Marson, L., MacGilchrist, A., Johannessen, I. (2013). High risk of cytomegalovirus infection following solid organ transplantation despite prophylactic therapy. *J Med Virol* 85, 893-898.
39. Heeger, P. S., Greenspan, N. S., Kuhlenschmidt, S., Dejelo, C., Hricik, D. E., Schulak, J. A., Tary-Lehmann, M. (1999). Pretransplant frequency of donor-specific, IFN-gamma-producing lymphocytes is a manifestation of immunologic memory and correlates with the risk of posttransplant rejection episodes. *J Immunol* 163, 2267-2275.
40. Ho, V. T., Soiffer, R. J. (2001). The history and future of T-cell depletion as graft-versus-host disease prophylaxis for allogeneic hematopoietic stem cell transplantation. *Blood* 98, 3192-3204.
41. Hoffman, F. M. (2005). Outcomes and complications after heart transplantation: a review. *J Cardiovasc Nurs* 20, S31-42.
42. Holl, E. K. (2013). Generation of bone marrow and fetal liver chimeric mice. *Methods Mol Biol* 1032, 315-321.
43. Hunt, S. A., Haddad, F. (2008). The changing face of heart transplantation. *J Am Coll Cardiol* 52, 587-598.
44. Hunziker, L., Recher, M., Macpherson, A. J., Ciurea, A., Freigang, S., Hengartner, H., Zinkernagel, R. M. (2003). Hypergammaglobulinemia and autoantibody induction mechanisms in viral infections. *Nat Immunol* 4, 343-349.

45. Ison, M. G., Nalesnik, M. A. (2011). An update on donor-derived disease transmission in organ transplantation. *Am J Transplant* 11, 1123-1130.
46. Jacobsohn, D. A., Vogelsang, G. B. (2007). Acute graft versus host disease. *Orphanet J Rare Dis* 2, 35.
47. Jagasia, M., Arora, M., Flowers, M. E., Chao, N. J., McCarthy, P. L., Cutler, C. S., Urbano-Ispizua, A., Pavletic, S. Z., Haagenson, M. D., Zhang, M. J., Antin, J. H., Bolwell, B. J., Bredeson, C., Cahn, J. Y., Cairo, M., Gale, R. P., Gupta, V., Lee, S. J., Litzow, M., Weisdorf, D. J., Horowitz, M. M., Hahn, T. (2012). Risk factors for acute GVHD and survival after hematopoietic cell transplantation. *Blood* 119, 296-307.
48. Jiang, H., Chess, L. (2009). How the immune system achieves self-nonself discrimination during adaptive immunity. *Adv Immunol* 102, 95-133.
49. Jung, S. H., Kim, J. J., Choo, S. J., Yun, T. J., Chung, C. H., Lee, J. W. (2011). Long-term mortality in adult orthotopic heart transplant recipients. *J Korean Med Sci* 26, 599-603.
50. Kissmeyer-Nielsen, F., Olsen, S., Petersen, V. P., Fjeldborg, O. (1966). Hyperacute rejection of kidney allografts, associated with pre-existing humoral antibodies against donor cells. *Lancet* 2, 662-665.
51. Klymenko, S. V., Belyi, D. A., Ross, J. R., Owzar, K., Jiang, C., Li, Z., J, N. M., A, N. K., Bebeshko, V. G., N, J. C. (2011). Hematopoietic cell infusion for the treatment of nuclear disaster victims: new data from the Chernobyl accident. *Int J Radiat Biol* 87, 846-850.
52. Knapp, S., Hennig, B. J., Frodsham, A. J., Zhang, L., Hellier, S., Wright, M., Goldin, R., Hill, A. V., Thomas, H. C., Thursz, M. R. (2003). Interleukin-10 promoter polymorphisms and the outcome of hepatitis C virus infection. *Immunogenetics* 55, 362-369.
53. Kroger, N., Zabelina, T., Kruger, W., Renges, H., Stute, N., Rischewski, J., Sonnenberg, S., Ayuk, F., Togel, F., Schade, U., Fiegel, H., Erttmann, R., Loliger, C., Zander, A. R. (2002). In vivo T cell depletion with pretransplant anti-thymocyte globulin reduces graft-versus-host disease without increasing relapse in good risk myeloid leukemia patients after stem cell transplantation from matched related donors. *Bone Marrow Transplant* 29, 683-689.
54. Kuhn, R., Lohler, J., Rennick, D., Rajewsky, K., Muller, W. (1993). Interleukin-10-deficient mice develop chronic enterocolitis. *Cell* 75, 263-274.

55. Lang, K. S., Georgiev, P., Recher, M., Navarini, A. A., Bergthaler, A., Heikenwalder, M., Harris, N. L., Junt, T., Odermatt, B., Clavien, P. A., Pircher, H., Akira, S., Hengartner, H., Zinkernagel, R. M. (2006). Immunoprivileged status of the liver is controlled by Toll-like receptor 3 signaling. *J Clin Invest* 116, 2456-2463.
56. Lang, K. S., Recher, M., Junt, T., Navarini, A. A., Harris, N. L., Freigang, S., Odermatt, B., Conrad, C., Ittner, L. M., Bauer, S., Luther, S. A., Uematsu, S., Akira, S., Hengartner, H., Zinkernagel, R. M. (2005). Toll-like receptor engagement converts T-cell autoreactivity into overt autoimmune disease. *Nat Med* 11, 138-145.
57. Lang, K. S., Recher, M., Navarini, A. A., Freigang, S., Harris, N. L., van den Broek, M., Odermatt, B., Hengartner, H., Zinkernagel, R. M. (2005). Requirement for neutralizing antibodies to control bone marrow transplantation-associated persistent viral infection and to reduce immunopathology. *J Immunol* 175, 5524-5531.
58. Lang, P. A., Recher, M., Haussinger, D., Lang, K. S. (2010). Genes determining the course of virus persistence in the liver: lessons from murine infection with lymphocytic choriomeningitis virus. *Cell Physiol Biochem* 26, 263-272.
59. Lehrer, R. I. (2004). Primate defensins. *Nat Rev Microbiol* 2, 727-738.
60. Ljungman, P. (2008). CMV infections after hematopoietic stem cell transplantation. *Bone Marrow Transplant* 42 Suppl 1, S70-S72.
61. Mielcarek, M., Martin, P. J., Leisenring, W., Flowers, M. E., Maloney, D. G., Sandmaier, B. M., Maris, M. B., Storb, R. (2003). Graft-versus-host disease after nonmyeloablative versus conventional hematopoietic stem cell transplantation. *Blood* 102, 756-762.
62. Mihatsch, M. J., Kyo, M., Morozumi, K., Yamaguchi, Y., Nickleleit, V., Ryffel, B. (1998). The side-effects of ciclosporine-A and tacrolimus. *Clin Nephrol* 49, 356-363.
63. Mohty, M., Labopin, M., Volin, L., Gratwohl, A., Socie, G., Esteve, J., Tabrizi, R., Nagler, A., Rocha, V., Acute Leukemia Working Party of, E. (2010). Reduced-intensity versus conventional myeloablative conditioning allogeneic stem cell transplantation for patients with acute lymphoblastic leukemia: a retrospective study from the European Group for Blood and Marrow Transplantation. *Blood* 116, 4439-4443.
64. Moore, K. W., de Waal Malefyt, R., Coffman, R. L., O'Garra, A. (2001). Interleukin-10 and the interleukin-10 receptor. *Annu Rev Immunol* 19, 683-765.

65. Moreno Brea, M. A., Mico, J. A. (2009). [TNF and cytokines and pain: Beyond the tisular inflammation]. *Reumatol Clin* 5 Suppl 2, 1-4.
66. Mori, S., Kocak, U., Shaw, J. L., Mullen, C. A. (2005). Augmentation of post transplant immunity: antigen encounter at the time of hematopoietic stem cell transplantation enhances antigen-specific donor T-cell responses in the post transplant repertoire. *Bone Marrow Transplant* 35, 793-801.
67. Nichols, W. G., Corey, L., Gooley, T., Davis, C., Boeckh, M. (2002). High risk of death due to bacterial and fungal infection among cytomegalovirus (CMV)-seronegative recipients of stem cell transplants from seropositive donors: evidence for indirect effects of primary CMV infection. *J Infect Dis* 185, 273-282.
68. Paladino, N., Fainboim, H., Theiler, G., Schroder, T., Munoz, A. E., Flores, A. C., Galdame, O., Fainboim, L. (2006). Gender susceptibility to chronic hepatitis C virus infection associated with interleukin 10 promoter polymorphism. *J Virol* 80, 9144-9150.
69. Persico, M., Capasso, M., Persico, E., Masarone, M., Renzo, A., Spano, D., Bruno, S., Iolascon, A. (2006). Interleukin-10 - 1082 GG polymorphism influences the occurrence and the clinical characteristics of hepatitis C virus infection. *J Hepatol* 45, 779-785.
70. Pietersma, F. L., van Dorp, S., Minnema, M. C., Kuball, J., Meijer, E., Schuurman, R., van Baarle, D. (2011). Influence of donor cytomegalovirus (CMV) status on severity of viral reactivation after allogeneic stem cell transplantation in CMV-seropositive recipients. *Clin Infect Dis* 52, e144-148.
71. Pircher, H., Burki, K., Lang, R., Hengartner, H., Zinkernagel, R. M. (1989). Tolerance induction in double specific T-cell receptor transgenic mice varies with antigen. *Nature* 342, 559-561.
72. Ramanan, P., Razonable, R. R. (2013). Cytomegalovirus infections in solid organ transplantation: a review. *Infect Chemother* 45, 260-271.
73. Razonable, R. R. (2011). Management of viral infections in solid organ transplant recipients. *Expert Rev Anti Infect Ther* 9, 685-700.
74. Razonable, R. R., Humar, A., Practice, A. S. T. I. D. C. o. (2013). Cytomegalovirus in solid organ transplantation. *Am J Transplant* 13 Suppl 4, 93-106.
75. Remberger, M., Mattsson, J., Ringden, O. (2001). Polyclonal anti-T-cell globulin as part of the preparative regimen for pediatric allogeneic stem-cell transplantation. *Pediatr Transplant* 5, 285-292.

76. Sarraj, B., Ye, J., Akl, A. I., Chen, G., Wang, J. J., Zhang, Z., Abadja, F., Abecassis, M., Miller, S. D., Kansas, G. S., Ansari, M. J. (2014). Impaired selectin-dependent leukocyte recruitment induces T-cell exhaustion and prevents chronic allograft vasculopathy and rejection. *Proc Natl Acad Sci U S A* 111, 12145-12150.
77. Schroder, K., Hertzog, P. J., Ravasi, T., Hume, D. A. (2004). Interferon-gamma: an overview of signals, mechanisms and functions. *J Leukoc Biol* 75, 163-189.
78. Schroeder, M. A., DiPersio, J. F. (2011). Mouse models of graft-versus-host disease: advances and limitations. *Dis Model Mech* 4, 318-333.
79. Sharpe, M., Mount, N. (2015). Genetically modified T cells in cancer therapy: opportunities and challenges. *Dis Model Mech* 8, 337-350.
80. Shin, H. D., Winkler, C., Stephens, J. C., Bream, J., Young, H., Goedert, J. J., O'Brien, T. R., Vlahov, D., Buchbinder, S., Giorgi, J., Rinaldo, C., Donfield, S., Willoughby, A., O'Brien, S. J., Smith, M. W. (2000). Genetic restriction of HIV-1 pathogenesis to AIDS by promoter alleles of IL10. *Proc Natl Acad Sci U S A* 97, 14467-14472.
81. Srinivasan, A., Burton, E. C., Kuehnert, M. J., Rupprecht, C., Sutker, W. L., Ksiazek, T. G., Paddock, C. D., Guarner, J., Shieh, W. J., Goldsmith, C., Hanlon, C. A., Zoretic, J., Fischbach, B., Niezgoda, M., El-Feky, W. H., Orciari, L., Sanchez, E. Q., Likos, A., Klintmalm, G. B., Cardo, D., LeDuc, J., Chamberland, M. E., Jernigan, D. B., Zaki, S. R., Rabies in Transplant Recipients Investigation, T. (2005). Transmission of rabies virus from an organ donor to four transplant recipients. *N Engl J Med* 352, 1103-1111.
82. Turner, D. (2004). The human leucocyte antigen (HLA) system. *Vox Sang* 87 Suppl1, 87-90.
83. Umemura, A., Morita, H., Li, X. C., Tahan, S., Monaco, A. P., Maki, T. (2001). Dissociation of hemopoietic chimerism and allograft tolerance after allogeneic bone marrow transplantation. *J Immunol* 167, 3043-3048.
84. van den Heuvel, H., Heidt, S., Roelen, D. L., Claas, F. H. (2015). T-cell alloreactivity and transplantation outcome: a budding role for heterologous immunity? *Curr Opin Organ Transplant* 20, 454-460.
85. Wherry, E. J. (2011). T cell exhaustion. *Nat Immunol* 12, 492-499.

86. Wherry, E. J., Ha, S. J., Kaech, S. M., Haining, W. N., Sarkar, S., Kalia, V., Subramaniam, S., Blattman, J. N., Barber, D. L., Ahmed, R. (2007). Molecular signature of CD8+ T cell exhaustion during chronic viral infection. *Immunity* 27, 670-684.
87. Winston, D. J., Vikram, H. R., Rabe, I. B., Dhillon, G., Mulligan, D., Hong, J. C., Busuttil, R. W., Nowicki, M. J., Mone, T., Civen, R., Tecle, S. A., Trivedi, K. K., Hocevar, S. N., West Nile Virus Transplant-Associated Transmission Investigation, T. (2014). Donor-derived West Nile virus infection in solid organ transplant recipients: report of four additional cases and review of clinical, diagnostic, and therapeutic features. *Transplantation* 97, 881-889.
88. Wu, K., Zhang, J., Fu, J., Wu, S., Philipp, T., Uwe, H., Kribben, A., Witzke, O. (2006). Novel technique for blood circuit reconstruction in mouse heart transplantation model. *Microsurgery* 26, 594-598.
89. Yi, J. S., Cox, M. A., Zajac, A. J. (2010). T-cell exhaustion: characteristics, causes and conversion. *Immunology* 129, 474-481.
90. Yusef, R. D., Edwards, L. B., Kucheryavaya, A. Y., Benden, C., Dipchand, A. I., Dobbels, F., Goldfarb, S. B., Levvey, B. J., Lund, L. H., Meiser, B., Stehlik, J., International Society for, H., Lung, T. (2014). The registry of the International Society for Heart and Lung Transplantation: thirty-first adult lung and heart-lung transplant report--2014; focus theme: retransplantation. *J Heart Lung Transplant* 33, 1009-1024.
91. Zinkernagel, R. M., Doherty, P. C. (1979). MHC-restricted cytotoxic T cells: studies on the biological role of polymorphic major transplantation antigens determining T-cell restriction-specificity, function, and responsiveness. *Adv Immunol* 27, 51-177.
92. Zinkernagel, R. M., Doherty, P. C. (1997). The discovery of MHC restriction. *Immunol Today* 18, 14-17.
93. Zinkernagel, R. M., Haenseler, E., Leist, T., Cerny, A., Hengartner, H., Althage, A. (1986). T cell-mediated hepatitis in mice infected with lymphocytic choriomeningitis virus. Liver cell destruction by H-2 class I-restricted virus-specific cytotoxic T cells as a physiological correlate of the 51Cr-release assay? *J Exp Med* 164, 1075-1092.
94. Zuo, Z., Wang, C., Carpenter, D., Okada, Y., Nicolaidou, E., Toyoda, M., Trento, A., Jordan, S. C. (2001). Prolongation of allograft survival with viral IL-10 transfection in a highly histoincompatible model of rat heart allograft rejection. *Transplantation* 71, 686-691.

7 Abbreviation

Ab	antibody
ALT	alanine transferase
AML	acute myeloid lymphoma
APC	antigen presenting cell
BMT	bone marrow transplantation
CAV	cardiac allograft vasculopathy
CD	cluster of differentiation
CFSE	carboxy-flourescein-succinimidyl-ester
CK-MB	creatinine kinase of type muscle brain
CMV	cytomegalovirus
DMEM	dulbecco`s modified eagle medium
DNA	deoxyribonucleic acid
ELISA	enzyme linked immunosorbent assay
FACS	fluorescence activated cell sorting
FCS	fetal calf serum
FITC	fluorescein-isothiocyanate
GvHD	graft versus host disease
GvL	graft versus leukemia
GP	glycoprotein
Gy	gray
HBV	hepatitis B virus
HCV	hepatitis C virus
HLA	human leukocyte antigen

HTX	heart transplantation
<i>i.p.</i>	intraperitoneal
<i>i.v.</i>	intravenous
IVC	inferior vena cava
IL	interleukin
IMDM	iscoves modified dulbeccos medium
IFN- γ	interferon gamma
LCMV	lymphocytic choriomeningitis virus
LDH	lactate dehydrogenase
LPS	lipopolysaccharide
MC57	mouse fibrosarcoma cell line
NAT	nucleic acid testing
PBS	phosphate buffered saline
PCR	polymerase chain reaction
PE	phycoerythrin
PeCy7	phycoerythrin-cyanin-7
PerCP	peridin-chlorophyll
PFU	plaque forming units
PSG	penicilin-streptomycin-glutamine
qRT-PCR	quantitative real time PCR
RIC	reduced intensity conditioning
SOT	solid organ transplantation
TCR	t cell receptor
TLR	toll-like receptor

Trop I	troponine I
WT	wild type

8 Tables and Figures

Fig. 1.1	Genetic organization of the major histocompatibility complex in human and in mouse.....	9
Fig. 1.2	T cell exhaustion and immunopathology.....	12
Fig. 3.1	LCMV-GP expression in <i>DEE</i> mice.....	28
Fig. 3.2	Presence of GP33 specific CD8 ⁺ T cells in thymus and blood.....	28
Fig. 3.3	Model of BMT.....	29
Fig. 3.4	Development of P14 T cells after BMT and measurement of ALT and LDH in sera.....	30
Fig. 3.5	P14 T cells generated in chimera mice got activated <i>ex-vivo</i>	31
Fig. 3.6	P14 T cell kinetic in blood of chimeric mice after mixed bone marrow transplantation.....	32
Fig. 3.7	Model of heterotopic heart transplantation.....	33
Fig. 3.8	Graft survival curve after HTX.....	33
Fig. 3.9	Tetramer staining of blood after HTX.....	34
Fig. 3.10	Experiment design.....	34
Fig. 3.11	Virus titers in LCMV donor hearts.....	35
Fig. 3.12	Virus titers in organs after LCMV-HTX.....	36
Fig. 3.13	CD8 ⁺ T cell kinetic and IFN- γ production in blood.....	37
Fig. 3.14	ALT and LDH measurement in serum after LPS challenge.....	38
Fig. 3.15	Experiment design.....	39
Fig. 3.16	Survival curve and CD8 ⁺ T cell kinetic after LCMV-HTX.....	40
Fig. 3.17	IL-10 and virus titers in plasma and blood after LCMV-HTX.....	40
Fig. 3.18	Analysis of heart and liver enzymes in sera after LCMV-HTX.....	41
Fig. 3.19	Experiment design.....	42

Fig. 3.20 Characterization of Memory T cells..... 43

Fig. 3.21 Virus control and transplant rejection in memory mice..... 43

9 Acknowledgment

I would like to thank everyone who supported me during my studies and my thesis.

First of all, I thank my parents, my sisters and my little brother who were and are always there for me and who encourage me in what I do and in what I am.

I thank my thesis supervisor, Prof. Dr. med. Karl Sebastian Lang, who was very patient and supported me and guided me through my doctoral medical thesis. I am very lucky that I had the opportunity to write my thesis in his lab. I learned and specialized in the vast field of research in immunology with his help.

Special thanks go to Konstanze Schättel who was always there for me, who understood me and who is the kindest and coolest person in the world.

I thank my colleagues Halime Kalkavan, Vikas Duhan, Vishal Khairnar, Namir Shaabani, Nadine Honke, Katja Merches, Piyush Sharma, Aleks Pandyra, Sarah-Kim Friedrich, Anna Höwner and Patricia Spieker for a great time. It was always fun working in the lab as a super team.

A big thanks is dedicated for Dr. Fu Jian, who performed the heart transplantation experiments. He is an awesome surgeon, very patient and kind. I thank Prof. Dr. med. Witzke and PD. Dr. med. Sebastian Dolff for the smooth cooperation.

Looking back, I am very grateful for the time that I had.

10 Curriculum vitae

Der Lebenslauf ist in der Online-Version aus Gründen des Datenschutzes nicht enthalten.